

Oreocharis corallodiscoides (Gesneriaceae, Trichosporeae, Didymocarpinae), a new species from Yunnan, southwest China

Xi Li¹, Qiu-Ping Wang¹, Feng Yang¹, Jing-Yi Ye¹, Huan-Chong Wang^{1,2}

¹ School of Ecology and Environmental Science, Yunnan University, Kunming 650091, China

² Herbarium of Yunnan University, Kunming 650091, Yunnan, China

Corresponding author: Huan-Chong Wang (hchwang@ynu.edu.cn)

Abstract

Oreocharis corallodiscoides, a new species of the tribe Trichosporeae (Gesneriaceae), is described and illustrated. This new species is characterized by its ovate, rounded or sometimes rhombic leaves densely covered with appressed hairs, yellow corolla, and the presence of two purple triangular appendages inside the base of the corolla tube. It is known only from the type locality, Luzhijiang dry-hot valley in Yimen county, central Yunnan, southwest China. A morphological comparison with its morphologically similar species (*O. agnesiae* and *O. longifolia*) is also presented.

Key words: Dry-hot valley, endemism, morphological comparison, taxonomy, xerophyte



Academic editor: John L Clark

Received: 5 February 2025

Accepted: 17 April 2025

Published: 13 May 2025

Citation: Li X, Wang Q-P, Yang F, Ye J-Y, Wang H-C (2025) *Oreocharis corallodiscoides* (Gesneriaceae, Trichosporeae, Didymocarpinae), a new species from Yunnan, southwest China. PhytoKeys 256: 73–81. <https://doi.org/10.3897/phytokeys.256.148644>

Copyright: © Xi Li et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The genus *Oreocharis* Benth. was first established by the British botanist G. Bentham (1876). For a long time, it was regarded as a medium-sized genus comprising approximately twenty to thirty species (Wang et al. 1998). Based on molecular data and morphological evaluation, Möller et al. (2011) expanded the delimitation of *Oreocharis* by incorporating ten small-sized or monotypic genera (e.g. *Ancylostemon* Craib, *Dayaoshania* W.T.Wang, *Isometrum* Craib, *Paraisometrum* W.T. Wang, *Thamnocharis* W.T.Wang, etc.) into a newly delimited *Oreocharis*. Additionally, several species previously placed in *Briggsia* Craib were transferred to *Oreocharis*, thereby expanding the genus to comprise approximately 90 species (Chen et al. 2014; Möller et al. 2014). The revised genus is characterized by rosette-forming herbs with spirally arranged leaves, scape inflorescences, and loculicidally dehiscent capsules (Möller et al. 2011). Beyond these shared traits, members of *Oreocharis* exhibit considerable variation in flower morphology (Möller et al. 2011). The treatment of Möller et al. (2011) is generally accepted by recent researchers, such as Fu et al. (2019), Tan et al. (2020), Yang et al. (2020), Li et al. (2023), Xie et al. (2024) and others. After the generic redefinition, about 70 species (Möller 2019; Cai et al. 2020; Chen et al. 2020; Yang et al. 2020) of *Oreocharis* have been described and officially published. Currently, the genus *Oreocharis* contains approximately 160 species (GRC 2025). This genus is mainly distributed in southern and southwestern

China, with a few species extending into or occurring in northern Vietnam, India, Bhutan, Japan, Thailand and Myanmar (Möller et al. 2011; Cai et al. 2020). Since 2011, approximately 70 new species of *Oreocharis* have been continuously discovered and published (e.g. Cai et al. 2019, 2020; Chen et al. 2020; Yang and Shi 2021; and summarized in Möller 2019; Wen et al. 2025). Most of these new species are from China, highlighting the need for continued investigation and study of *Oreocharis* diversity in China.

During botanical fieldwork in the Luzhijiang valley, Yimen county, Yunnan province, southwest China, in September 2015, we discovered and collected an unknown species of Gesneriaceae. After a detailed comparison with morphologically similar species, we identified it as a member of *Oreocharis* based on its inflated corolla tube, widely open corolla, and yellow floral coloration—features shared with *O. longifolia* (the type species of the genus *Briggsia*, now placed in *Oreocharis*), and this plant represents a distinct new species.

Materials and methods

The study followed standard practices in plant taxonomic surveys and herbarium research. Morphological analyses of the new species were examined through observations of living plants and herbarium specimens. Observation of live plants was carried out on the field population, with approximately 30 individuals directly analyzed and studied. Meanwhile, fresh flowers were brought to the laboratory for dissection and observation. Digital images of type specimens of the genus *Oreocharis*, available at the JSTOR Global Plants (<http://plants.jstor.org/>), were thoroughly examined. Additionally, collections (actual specimens or digital photos) housed in the herbaria BM, E, NE, GH, KUN, and YUKU (following acronyms standardized by Thiers (2025) [updated continuously]) were carefully reviewed and compared to the new species. The pertinent taxonomic literature (Wang et al. 1990, 1998; Wei et al. 2010; Möller 2019; Wen et al. 2025; GRC 2025) was consulted extensively to ensure that no known species were overlooked.

Taxonomic treatment

***Oreocharis corallodiscoides* Huan C. Wang & Xi Li, sp. nov.**

urn:lsid:ipni.org:names:77361684-1

Figs 1–3

Diagnosis. *Oreocharis corallodiscoides* is similar to *O. agnesiae* (Forrest ex W. W. Sm.) Mich. Möller et W. H. Chen in texture and indumentum of the leaves, as well as in flower morphology and size, but can be clearly distinguished from the latter by its leaves ovate, rounded or sometimes rhombic (vs. ovate to oblong), with a length-to-width ratio of 1–1.5 (vs. 2.2–2.7), corollas yellow (vs. purple-red), and the presence (vs. absence) of two purple and triangular appendages inside the base of the corolla tube. Additionally, while *O. corallodiscoides* shares the inflated yellow corolla tube with *O. longifolia*, it differs by its ovate-rotund leaves (vs. lanceolate-oblong), shorter petioles (0.5–2 cm vs. 3.5–5 cm), crenate leaf margins (vs. serrate), glandular pistils (vs. glabrous), and the presence of appendages with thickened filaments.




Figure 1. *Oreocharis corallodiscoides* sp. nov. **A** habit **B** adaxial surface of leaf **C** abaxial surface of leaf **D** flower (side view) **E** style, stamens and two triangular purple appendages **F** immature capsule.

Type. CHINA • Yunnan province, Yimen county, Luzhi town, near Xiaoluzhi village, 24°40'N, 101°57'E, 1300–1400 m a.s.l., 25th Sept. 2015, H. C. Wang et al. YM239 (Holotype: YUKU [2074890]!; isotypes: KUN, YUKU).

Description. *Herbs perennial.* Rhizome inconspicuous, with numerous fibrous roots. **Leaves** 9–18 in basal rosette, petiolate; **petiole** complanate,

0.5–2 cm long, both surfaces with densely appressed multicellular hairs; **leaf blade** ovate, rotund, or sometimes rhombic, $2.2\text{--}3.5 \times 1.1\text{--}2.2$ cm, coriaceous when dried; apex rounded, base broadly cuneate to rounded, margins crenate, adaxial surfaces densely white to gray strigose, abaxial with densely appressed multicellular hairs; **lateral veins** 4 or 5 on each side of midrib, adaxially inconspicuous, abaxially conspicuous. **Cymes** 1–2 (–4), scapiform, 1-flowered; **peduncles** 4–7 cm long, with dense glandular hairs; **bracts** 2, inserted above middle of cymes, elliptic to lanceolate, $2.0\text{--}4.0 \times \text{ca. } 1.0$ mm, with entire margins, outside with glandular hair, inside nearly glabrous; **pedicels** 2–3 cm long, with dense glandular hairs. **Calyx** 5-parted to near base, narrowly triangular, apex acute, margin entire, outside with glandular hair, inside nearly glabrous, sepals unequal, 4–6 mm long. **Corolla** yellow, not spotted or striped, 2.0–3.5 cm long, ca. 1–1.2 cm wide at mouth, outside covered with glandular hairs, glabrous inside; **tube** 1.5–2.5 cm long, inflated, infundibuliform and slightly curved downwards, two triangular purple appendages attached to base of tube, ca. 3 mm long, 2 mm wide at the base, apex acute; **limb** 5–6-lobed, distinctly 2-lipped, adaxial lip 4–6 mm long, unlobed, apex with 3 teeth, abaxial lip 7–8 mm long, 3-lobed, lobes wide triangular, 3–4 mm wide, apex acuminate; **disk** ring-like, ca. 1 mm in height; **fertile stamens** 4, adnate to corolla base, coherent in pairs, included, filaments ca. 2 cm in length; **staminode** 1, ca. 1 cm in length, with glandular hairs, degenerate anthers reniform. **Pistil** with glandular hairs, 2–2.5 cm long, ovary long cylindrical, light yellow, 1-loculed, style 1–2 mm long, stigma bilobed. **Capsule** oblanceolate-oblong, 4–6 cm long, commonly one side of the capsule dehiscing first.

Phenology. *Oreocharis corallodiscoides* was observed flowering from August to September, and fruiting in October.

Etymology. The specific epithet *corallodiscoides* is derived from the generic name *Corallodiscus* Batalin (Gesneriaceae) by adding the suffix “-oides”, reflecting the leaf morphological similarity of this new species to some members of the genus *Corallodiscus*, such as *C. lanuginosus* (Wall. ex A. DC.) B. L. Burtt; particularly, its sometimes rhombic leaves are rare in the genus *Oreocharis*.

Vernacular name. Shan hu ye fo du ju tai (珊瑚叶佛肚菖苔) (Chinese).

Distribution and ecology. *Oreocharis corallodiscoides* is known only from Xiaoluzhi village in the Luzhijiang valley, Yimen county, Yunnan province, southwest China. Its habitat is characterized by seasonally hot and arid conditions, occurring on dry slopes within the valley (Fig. 3). The species is typically found in limestone grasslands at elevations ranging from 1,300 to 1,400 meters, where water availability is limited. Additionally, *Oreocharis corallodiscoides* often grows together with *Corallodiscus lanuginosus* (Wall. ex A. DC.) B. L. Burtt, and interestingly, these two species of different genera have very similar leaf shapes and indumentum (Fig. 3B).

Conservation status. *Oreocharis corallodiscoides* is a rare species with a restricted geographical range and a small population size. It is currently known only from the upstream region of the Luzhijiang river in Yimen county, an area that lies outside any officially protected zone. Based on the IUCN Red List Categories and Criteria (Version 16) (IUCN, 2024), the species meets Criterion D (very small or restricted population), with total mature individuals fewer than 250 mature individuals. Thus, this newly described species is preliminarily assessed as Endangered (EN).




Figure 2. *Oreocharis corallodiscoides* sp. nov. **A, B** habit **C** flower in front view **D** flower in side view, pedicel **E** adaxial surface of leaf **F** abaxial surface of leaf **G** pistil and stamens **H** pistil, stamens and two triangular purple glandular appendages (at the arrow) **I** immature capsule **J** mature capsule.



Figure 3. Habitat of *Oreocharis corallodiscoides* sp. nov. **A** overview of the Luzhijiang valley showing the type locality (indicated by red arrow) **B** close-up of the fruiting habitat (*Oreocharis corallodiscoides* and *Corallodiscus lanuginosus* are indicated by red and blue arrows respectively) **C** close-up of the xeric habitat during the dry season.

Discussion

Given its habitat and morphological characteristics, *Oreocharis corallodiscoides* appears to be a xerophytic plant well adapted to arid environments. This new species has leathery, thick-textured leaves densely covered with an indumentum and characterized by a relatively small leaf area. Unlike many congeners that grow on damp, shady rocks or cliffs in forested or rocky areas (Wei et al. 2010), *O. corallodiscoides* is found on open, dry, rocky slopes within dry-hot valleys. Additionally, the presence of two purple, triangular appendages at the base of the corolla tube (Fig. 2H) represents a character unique to this species, as it has not been observed in other members of the genus by us or documented in existing literature.

Morphologically, *Oreocharis corallodiscoides* resembles *O. agnesiae*, sharing similar leaf indumentum and flower morphology. However, these two species show significant differences in corolla color: *O. corallodiscoides* has a uniformly yellow corolla, whereas the corolla of *O. agnesiae* is purple. Additionally, their leaf morphology also differs markedly. Geographically, *O. corallodiscoides* has a distinct and non-overlapping distribution range with *O. agnesiae* (Fig. 4). *Oreocharis corallodiscoides* is restricted to the Luzhijiang valley, an upstream

tributary of the Yuanjiang river, at elevations of 1,300–1,400 m in central Yunnan. In contrast, *O. agnesiae* is distributed in the valleys of the Jinsha river and its tributaries at elevations of 1,900–3,000 m in northwestern Yunnan and southwestern Sichuan (Fig. 4). *Oreocharis corallodiscoides* is morphologically similar to *O. longifolia* (Craib) Mich. Möller et A. Weber in sharing inflated corolla tube and yellow color. However, *O. corallodiscoides* obviously differs from *O. longifolia* by its leaf blade shape (ovate to rotund vs. lanceolate to oblong), petiole length (0.5–2 vs. 3.5–5 cm), leaf length (2.2–3.5 vs. 8–23 cm), leaf margin (crenate vs. serrate), indumentum of pistil (glandular hair vs. glabrous), appendages (presence vs. absence) and filaments (thickened vs. not thickened).

Additional specimens examined (Paratypes). CHINA • Yunnan: Yimen County, Luzhi town, Xiaoluzhi village, elev. ca. 1340 m, 22 Oct. 2015, H. C. Wang et al. YM324 (YUKU).




Figure 4. Geographic distribution of *Oreocharis corallodiscoides* and its morphologically similar species *O. agnesiae*.

Acknowledgements

We sincerely thank Dr. Michael Moeller for sharing his taxonomic expertise and constructive feedback.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was supported by the National Natural Science Foundation of China (Grant No. 31960040) and the Research on the Persistent Environmental Impacts of Hydropower Development in the Wunonglong to Nan'a River Mouth Section of the Lancang River Mainstream (HY2021/S17-1).

Author contributions

Conceptualization: HCW, XL. Investigation: HCW, FY, XL, QPW, YJY. Resources: XL. Visualization: QPW. Writing - original draft: XL. Writing - review and editing: XL, FY, HCW.

Author ORCIDs

Xi Li  <https://orcid.org/0009-0003-8498-7809>

Feng Yang  <https://orcid.org/0000-0003-2076-0130>

Jing-Yi Ye  <https://orcid.org/0000-0002-3737-0115>

Huan-Chong Wang  <https://orcid.org/0000-0001-8562-8849>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bentham G (1876) Gesneriaceae. In: Bentham G, Hooker JD (Eds) *Genera Plantarum* 2 (2). Lovell Reeve & Co., London, 990–1025.
- Cai L, Guo Y, Zhang RM, Dao ZL, Wen F (2019) *Oreocharis panzhouensis* (Gesneriaceae), a new species from karst regions in Guizhou, China. *Phytotaxa* 393(3): 287–291. <https://doi.org/10.11646/phytotaxa.393.3.5>
- Cai L, Huang ZJ, Wen F, Dao ZL (2020) Two new species of *Oreocharis* (Gesneriaceae) from karst regions in Yunnan and notes on *O. tetraptera* and *O. brachypoda* from China. *PhytoKeys* 162: 1–12. <https://doi.org/10.3897/phytokeys.162.52174>
- Chen WH, Shui YM, Möller M (2014) Two new combinations in *Oreocharis* Benth. (Gesneriaceae) from China. *Candollea* 69(2): 179–182. <https://doi.org/10.15553/c2014v692a10>
- Chen WH, Zhang YM, He DM, Li YL, Shui YM (2020) Four new species of *Oreocharis* (Gesneriaceae) in Yunnan province, China. *PhytoKeys* 157: 83–99. <https://doi.org/10.3897/phytokeys.157.32284>
- Fu Q, Xia Y, Guo Y, Huang R, Wang YQ (2019) *Oreocharis odontopetala*, a new species of Gesneriaceae from Guizhou, China. *PhytoKeys* 124: 1–9. <https://doi.org/10.3897/phytokeys.124.34609>
- GRC (2025) [continuously updated] Gesneriaceae Resource Centre. Royal Botanic Garden Edinburgh. <https://padme.rbge.org.uk/GRC> [Retrieved/Accessed: 1 April 2025]
- IUCN (2024) IUCN Red List Categories and Criteria (Version 16). International Union for Conservation of Nature. <https://www.iucnredlist.org/resources/redlistguidelines> [Accessed: 1 April 2025]
- Li RF, Le XG, Xu L, Maciejewski S, Chen B, Wen F (2023) *Oreocharis yangjifengensis* (Gesneriaceae), a new species from Yangjifeng National Nature Reserve of Yingtang

- City, Jiangxi Province, China. *Phytotaxa* 583(2): 213–218. <https://doi.org/10.11646/phytotaxa.583.2.10>
- Möller M (2019) Species discovery in time: An example from Gesneriaceae in China. *Guangxi Sciences* 26(1): 116. <https://doi.org/10.13656/j.cnki.gxkx.20190307.002>
- Möller M, Middleton D, Nishii K, Wei YG, Sontag S, Weber A (2011) A new delineation for *Oreocharis* incorporating an additional ten genera of Chinese Gesneriaceae. *Phytotaxa* 23(1): 1–36. <https://doi.org/10.11646/phytotaxa.23.1.1>
- Möller M, Chen WH, Shui YM, Atkins H, Middleton DJ (2014) A new genus of Gesneriaceae in China and the transfer of *Briggsia* species to other genera. *Gardens' Bulletin* (Singapore) 66: 195–205.
- Tan K, Lu T, Ren M-X (2020) Biogeography and evolution of Asian Gesneriaceae based on updated taxonomy. *PhytoKeys* 157: 7–26. <https://doi.org/10.3897/phytokeys.157.34032>
- Thiers B (2025) [continuously updated] Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden. <http://sweetgum.nybg.org/science/ih/> [Retrieved/Accessed: 1 April 2025]
- Wang WT, Pan KY, Li ZY (1990) Gesneriaceae. In: Wang WT (Ed.) *Flora Reipublicae Popularis Sinicae* (Vol. 69). Science Press, Beijing, 125–581.
- Wang WT, Pan KY, Li ZY, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) *Flora of China* (Vol. 18). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 244–401.
- Wei YG, Wen F, Möller M, Monro A, Zhang Q, Gao Q, Mou HF, Zhong SH, Cui C (2010) *Gesneriaceae of South China*. Guangxi Science and Technology Publishing House, Nanning, Guangxi, 777 pp.
- Wen F, Wei YG, Fu LF, Xin ZY, Ge YZ (2025) The Checklist of Gesneriaceae in China. <http://gccc.gxib.cn/cn/about-68.aspx> [Retrieved/Accessed: 1 April 2025]
- Xie Z, Peng NN, Zhang M, Ding GE, Wen F, Kong HH (2024) *Oreocharis scutifolia* (Gesneriaceae), a peltate-leaved new species from the dry-hot valley of the Jinsha River Basin, Yunnan, China. *Ecology and Evolution* 14(10): e70442. <https://doi.org/10.1002/ece3.70442>
- Yang LH, Shi XZ (2021) *Oreocharis reticuliflora* (Gesneriaceae), a new species from southeastern Sichuan, China. *Nordic Journal of Botany* 39(11): e03322. <https://doi.org/10.1111/njb.03322>
- Yang LH, Wen F, Kong HH, Sun ZX, Su LY, Kang M (2020) Two new combinations in *Oreocharis* (Gesneriaceae) based on morphological, molecular, and cytological evidence. *PhytoKeys* 157: 43–58. <https://doi.org/10.3897/phytokeys.157.32609>